

June 2025

2024 Minnesota Air Monitoring Data Report

2026 Minnesota Air Monitoring Network Plan

Authors

Kellie Gavin
Sanna Mairet
Margaret McCourtney
Tsheena Singh
Maria Takahashi
Nicholas Witcraft

Contributors/acknowledgements

Kurt Anderson
Yong Cai
Jerrod Eppen
Jeff Laren
Chloe Meyer
Nate Niebeling
Jacob Nelson
Binh Nguyen
Ashley Olson
Luke Salisbury
Mike Schneider
Sydney Schultz
Kristin Siewert
Joseph Smith
Michael Smith
Owen Seltz
Daniel Steltz
Matthew Taraldsen
Eric Wilcox Freeburg
David Wischnack

Editing

Jennifer Holstad

Minnesota Pollution Control Agency

520 Lafayette Road North | Saint Paul, MN 55155-4194 |

651-296-6300 | 800-657-3864 | Or use your preferred relay service | Info.pca@state.mn.us

This report is available in alternative formats upon request, and online at www.pca.state.mn.us.

Document number: aq10-26d

Table of contents

List of tables	1
List of figures.....	1
Acronyms, abbreviations, and definitions	2
Introduction	4
1. Pollutants with standards (criteria pollutants)	4
1.1. Air quality design values.....	4
1.2. National Ambient Air Quality Standards	4
1.3. Particulate Matter	5
1.3.1. PM _{2.5} - Fine particulate matter	6
1.3.2. PM ₁₀	10
1.3.3. Total Suspended Particulate matter (TSP)	12
1.4. Lead (Pb).....	14
1.5. Ozone (O ₃).....	16
1.6. Nitrogen Dioxide (NO ₂)	18
1.7. Carbon monoxide (CO)	20
1.8. Sulfur dioxide (SO ₂).....	22
2. AQI.....	23
3. Air Toxics	28
4. Visibility.....	28

List of tables

Table 1. National Ambient Air Quality Standards (NAAQS).....	5
Table 2. 2024 PM ₁₀ estimated exceedances.....	11
Table 3. Minnesota Ambient Air Quality Standards for TSP	12

List of figures

Figure 1. 2024 PM _{2.5} monitoring sites in Minnesota.	6
Figure 2. 2024 PM _{2.5} design values compared to the annual NAAQS.....	7
Figure 3. 2024 PM _{2.5} design values compared to the 24-hour NAAQS.....	8
Figure 4. Average PM _{2.5} concentrations per hour of the day at several Minnesota sites, May 2024	9
Figure 5. Daily average of hourly PM _{2.5} concentrations at several Minnesota sites, weekdays in March 2024	9
Figure 6. 2024 PM ₁₀ monitoring sites in Minnesota.	10
Figure 7. 2024 PM ₁₀ background concentrations compared to the daily NAAQS.	11
Figure 8. 2024 TSP monitoring sites in Minnesota	12
Figure 9. 2024 TSP concentrations compared to the annual primary and secondary MAAQS.	13
Figure 10. 2024 TSP concentrations compared to the 24-hour primary and secondary MAAQS.	13
Figure 11. 2024 lead monitoring sites in Minnesota.....	14
Figure 12. 2024 lead design values compared to the 3-month NAAQS.....	15

Figure 13. 2024 ozone monitoring sites in Minnesota.	16
Figure 14. 2024 ozone design values compared to the 8-hour average NAAQS.	17
Figure 15. 2024 NO _x monitoring sites in Minnesota.....	18
Figure 16. 2024 NO ₂ design values compared to the annual NAAQS.	19
Figure 17. 2024 NO ₂ design values compared to the 1-hour NAAQS.	19
Figure 18. 2024 CO monitoring sites in Minnesota.	20
Figure 19. 2024 design values for CO compared to the 8-hour NAAQS.	21
Figure 20. 2024 CO design values compared to the 1-hour average NAAQS.	21
Figure 21. 2024 SO ₂ monitoring sites in Minnesota.	22
Figure 22. 2024 SO ₂ design values compared to the 1-hour NAAQS.....	22
Figure 23. The Air Quality Index categories and respective levels of health concern (from https://airnow.gov).	23
Figure 24. Annual statewide PM _{2.5} AQI category counts, 2001-2024.....	23
Figure 25. 2024 daily observed AQI category counts for PM _{2.5} in Minnesota.....	24
Figure 26. Peak daily PM _{2.5} AQI for Minnesota in 2024.....	25
Figure 27. Annual statewide ozone AQI category counts, 2001-2024.....	25
Figure 28. 2024 daily observed AQI category counts for ozone in Minnesota (during the official ozone season from March 1 to October 31).	26
Figure 29. Peak daily ozone AQI for Minnesota in 2024.....	26
Figure 30. Total count and causes of air quality alert forecasts issued in Minnesota counties, 2010-2024.	27
Figure 31. Total count and causes of air quality forecast alert forecast days issued in Minnesota counties, 2010-2024.	27
Figure 32. Class I areas in Minnesota impacted by regional haze.	29
Figure 33. Visibility progress measured at Boundary Waters and Voyageurs (2004-2023) and the estimated interim progress goals for 2018 and 2028.	29

Acronyms, abbreviations, and definitions

AIRNow – air quality forecasting program

Air toxics – suite of parameters that includes VOCs, carbonyls, and metals

AQI – Air Quality Index

AQS – Air Quality System: EPA's repository of ambient air quality data

CAA – Clean Air Act

CAS – Chemical Abstracts Service

CFR – Code of Federal Regulations

Class I area – remote area with pristine air quality

CO – carbon monoxide

Criteria pollutants – the six pollutants regulated by the 1970 Clean Air Act (particulate matter, ozone, nitrogen dioxide, sulfur dioxide, carbon monoxide, and lead)

IMP – Chemical Speciation Network

Design Value – a statistic that describes the air quality status of a given location relative to the level of the National Ambient Air Quality Standards (NAAQS)

EPA – U.S. Environmental Protection Agency

Exceptional Events – unusual or naturally occurring events that can affect air quality but are not reasonably controllable using techniques that tribal, state or local air agencies may implement in order to attain and maintain the National Ambient Air Quality Standards.

FEM – Federal Equivalent Method

FRM – Federal Reference Method
IMPROVE – Interagency Monitoring of Protected Visual Environments
MAAQS – Minnesota Ambient Air Quality Standard
MDH – Minnesota Department of Health
MDN – Mercury Deposition Network
µg/m³ (ug/m³) – micrograms per cubic meter
MOA – Memorandum of Agreement
MPCA – Minnesota Pollution Control Agency
MSA – Metropolitan Statistical Area
NAAQS – National Ambient Air Quality Standard
NADP – National Atmospheric Deposition Program
NCore – National Core Monitoring Network
NDDN – National Dry Deposition Network
NO – nitric oxide
NO₂ – nitrogen dioxide
NO_x – oxides of nitrogen
NO_y – total reactive nitrogen
NTN – National Trends Network
O₃ – ozone
PAMS – Photochemical Assessment Monitoring Stations
Pb – lead
PM_{2.5} – particulate matter less than 2.5 microns in diameter (fine particulate matter)
PM_{10-2.5} – particulate matter between 2.5 and 10 microns in diameter (coarse particulate matter)
PM₁₀ – particulate matter less than 10 microns in diameter
ppb – parts per billion by volume
ppm – parts per million by volume
PQAO – Primary Quality Assurance Organization
Primary Standard - NAAQS set to protect public health
QAPP – Quality Assurance Project Plans
QA/QC – Quality Assurance/Quality Control
QMP – Quality Management Plan
Secondary Standard - NAAQS set to protect the environment and public welfare (i.e. visibility, crops, animals, vegetation, and buildings)
SIP – State Implementation Plan
SLAMS – State and Local Air Monitoring Stations
SO₂ – sulfur dioxide
SPM – special purpose monitoring
TPY – tons per year
TRS – total reduced sulfur
TSP – total suspended particulate matter
U of M – University of Minnesota
UFP – ultrafine particles (particulate matter less than 0.1 microns in diameter)
USDA – United States Department of Agriculture
USDOI – United States Department of the Interior
USG – unhealthy for sensitive groups
VOC – volatile organic compound

Introduction

The Minnesota Pollution Control Agency's (MPCA) air quality data are used to determine compliance with NAAQS and Minnesota Ambient Air Quality Standards (MAAQS). In 1970, the Clean Air Act (CAA) established NAAQS for six pollutants known to cause harm to human health and the environment. The CAA requires the MPCA to monitor these pollutants, called criteria pollutants, and report the findings to the U.S. Environmental Protection Agency (EPA). The criteria pollutants are particulate matter, lead, ozone, nitrogen dioxide, sulfur dioxide, and carbon monoxide. The MPCA monitors criteria pollutants to comply with the CAA. All criteria pollutants also have MAAQS. In addition, there are also MAAQS for total suspended particulate matter (TSP) and H₂S.

The MPCA also monitors Minnesota's air for other pollutants, called air toxics. Air toxics include a wide range of chemicals that are known or suspected to affect human health. These pollutants do not have federal standards; however, levels found in Minnesota are compared to health benchmarks established by the Minnesota Department of Health (MDH), the EPA, and the State of California.

More information including current air quality, forecasts, tools to explore data from our monitoring network, and this plan can be found on the [MPCA's Air Quality website \(https://www.pca.state.mn.us/air\)](https://www.pca.state.mn.us/air).

1. Pollutants with standards (criteria pollutants)

In 1970, the CAA authorized the EPA to establish standards for six pollutants known to cause harm to human health and the environment; these were given the name criteria pollutants. The criteria pollutants are particulate matter (currently PM_{2.5} and PM₁₀), lead (Pb), ozone (O₃), nitrogen dioxide (NO₂), sulfur dioxide (SO₂), and carbon monoxide (CO).

1.1. Air quality design values

A design value is a statistic that describes the air quality status of a given location relative to the level of the National Ambient Air Quality Standards (NAAQS). Design values are defined to be consistent with the individual NAAQS as described in [40 CFR Part 50 \(https://www.ecfr.gov/current/title-40/chapter-1/subchapter-C/part-50?toc=1\)](https://www.ecfr.gov/current/title-40/chapter-1/subchapter-C/part-50?toc=1). They are typically used to designate and classify nonattainment areas, as well as to assess progress towards meeting the NAAQS. Areas of the country where air pollution levels persistently exceed the national ambient air quality standards may be designated "nonattainment." The [EPA Green Book \(https://www.epa.gov/green-book\)](https://www.epa.gov/green-book) provides current information on nonattainment areas by state and pollutant.

Design values are computed and published annually by EPA's Office of Air Quality Planning and Standards and reviewed in conjunction with the EPA Regional Offices. More information can be found on [EPA's Design Values webpage \(https://www.epa.gov/air-trends/air-quality-design-values\)](https://www.epa.gov/air-trends/air-quality-design-values).

1.2. National Ambient Air Quality Standards

For each of these pollutants, the EPA has developed national ambient air monitoring quality standards (NAAQS). Primary standards are set to protect public health, while secondary standards are set to protect the environment and public welfare (i.e. visibility, crops, animals, vegetation, and buildings).

The CAA requires the EPA to review the scientific basis of these standards every five years to ensure they are protective of public health and the environment. The latest updates to the standards can be

found at [EPA's NAAQS webpage \(https://www.epa.gov/criteria-air-pollutants/naqs-table\)](https://www.epa.gov/criteria-air-pollutants/naqs-table), also shown in Table 1.

Table 1. National Ambient Air Quality Standards (NAAQS).

Pollutant [final rule citation]	Primary/ Secondary	Averaging time	Level	Form	
Carbon monoxide (CO) [76 FR 54294, Aug 31, 2011]	Primary	8 hours	9 ppm	Not to be exceeded more than once per year	
		1 hour	35 ppm		
Lead (Pb) [73 FR 66964, Nov 12, 2008]	Primary and secondary	Rolling 3-month average	0.15 µg/m ³ ⁽¹⁾	Not to be exceeded	
Nitrogen dioxide (NO₂) [75 FR 6474, Feb 9, 2010] [77 FR 20218, April 3, 2012]	Primary	1 hour	100 ppb	98 th percentile of 1-hour daily maximum concentrations, averaged over 3 years	
	Primary and secondary	1 year	53 ppb ⁽²⁾	Annual Mean	
Ozone (O₃) [80 FR 65292, Oct 26, 2015]	Primary and secondary	8 hours	0.070 ppm ⁽³⁾	Annual fourth-highest daily maximum 8-hour concentration, averaged over 3 years	
Particle Pollution (PM) 78 FR 3086, Jan 15, 2013	PM _{2.5}	Primary	1 year	9.0 µg/m ³	Annual mean, averaged over 3 years
		Secondary	1 year	15.0 µg/m ³	Annual mean, averaged over 3 years
		Primary and secondary	24 hours	35 µg/m ³	98 th percentile, averaged over 3 years
	PM ₁₀	Primary and secondary	24 hours	150 µg/m ³	Not to be exceeded more than once per year on average over 3 years
Sulfur dioxide (SO₂) [75 FR 35520, Jun 22, 2010] [84 FR 9866, April 17, 2019]	Primary	1 hour	75 ppb ⁽⁴⁾	99 th percentile of 1-hour daily maximum concentrations, averaged over 3 years	
	Secondary	3 hours	0.5 ppm	Not to be exceeded more than once per year	

(1) In areas designated nonattainment for the Pb standards prior to the promulgation of the current (2008) standards, and for which implementation plans to attain or maintain the current (2008) standards have not been submitted and approved, the previous standards (1.5 µg/m³ as a calendar quarter average) also remain in effect.

(2) The level of the annual NO₂ standard is 0.053 ppm. It is shown here in terms of ppb for the purposes of clearer comparison to the 1-hour standard level.

(3) Final rule signed October 1, 2015, and effective December 28, 2015. The previous (2008) O₃ standards additionally remain in effect in some areas. Revocation of the previous (2008) O₃ standards and transitioning to the current (2015) standards will be addressed in the implementation rule for the current standards.

(4) The previous SO₂ standards (0.14 ppm 24-hour and 0.03 ppm annual) will additionally remain in effect in certain areas: (1) any area for which it is not yet 1 year since the effective date of designation under the current (2010) standards, and (2) any area for which an implementation plan providing for attainment of the current (2010) standard has not been submitted and approved and which is designated nonattainment under the previous SO₂ standards or is not meeting the requirements of a SIP call under the previous SO₂ standards (40 CFR 50.4(3)). A SIP call is an EPA action requiring a state to resubmit all or part of its State Implementation Plan to demonstrate attainment of the required NAAQS.


1.3. Particulate Matter

The MPCA monitors four different sizes of particulates: fine particulate matter (PM_{2.5}), coarse particulate matter (PM_{10-2.5}), PM₁₀, and TSP. PM_{2.5} and PM₁₀ are regulated by the NAAQS and TSP is regulated by the MAAQS. There are currently no state or federal air quality standards for PM_{10-2.5}.

1.3.1. PM_{2.5} - Fine particulate matter

There were 26 FRM and FEM PM_{2.5} sites in Minnesota in 2024, 11 in the Twin Cities metropolitan area.

Figure 1. 2024 PM_{2.5} monitoring sites in Minnesota.




PM_{2.5} Regulatory network overview

The PM_{2.5} regulatory network includes FRM and FEM monitors. Currently, the MPCA operates FRM samplers at five sites and FEM monitors at 24 sites (Figure 1). Since PM_{2.5} NAAQS are based on three years of data and Mankato (5510) and Minneapolis (1904) were established in 2024; those sites are not included in the NAAQS charts below.

On February 7, 2024, EPA strengthened the NAAQS for Particulate Matter. EPA is changing the level of the primary (health-based) annual PM_{2.5} standard from 12.0 micrograms per cubic meter to 9.0 micrograms per cubic meter to provide increased public health protection, consistent with the available health science. EPA is not changing the current secondary annual PM_{2.5} standard nor the daily PM_{2.5} standard. For more information, visit <https://www.epa.gov/pm-pollution/final-reconsideration-national-ambient-air-quality-standards-particulate-matter-pm>.

A given monitoring site meets the annual PM_{2.5} NAAQS if the three-year average of the annual average PM_{2.5} concentration is less than or equal to 9 µg/m³. Results from PM_{2.5} monitors in 2022-2024 show Minnesota annual average PM_{2.5} concentrations ranged from 3.1 µg/m³ in Grand Portage (7810) to 8.4 µg/m³ in Minneapolis (0963); all below the current annual standard of 9 µg/m³ (Figure 2). 2024 design values have decreased relative to previous years because the calculations no longer include data from the 2021 wildfire smoke season. However, 2023 wildfire smoke values are still included in the 2024 design value calculations.


Figure 2. 2024 PM_{2.5} design values compared to the annual NAAQS



*Site did not meet data completeness criteria

A site meets the 24-hour NAAQS if the 98th percentile of the 24-hour PM_{2.5} concentrations in a year, averaged over three years, is less than or equal to 35 µg/m³. Results from FEM monitors in 2022-2024 show the 98th percentile of the daily PM_{2.5} averages in Minnesota ranged from 14 µg/m³ in Grand Portage (8710) to 27 µg/m³ in Marshall (4210) and Minneapolis (0963); all below the 24-hour standard of 35 µg/m³ (Figure 3).

Figure 3. 2024 PM_{2.5} design values compared to the 24-hour NAAQS.




*Site did not meet data completeness criteria

PM_{2.5} Continuous network

The MPCA currently supports 24 FEM PM_{2.5} sites in Minnesota. In addition to providing NAAQS comparable data, the PM_{2.5} continuous (FEM) data provide two key types of information that are not available from the FRM network. First, continuous data capture high concentration days that might be missed in the one-in-three-day FRM sampling schedule. Second, daily monitoring allows for temporal contrasts between sites on an ongoing basis, providing better, more informative comparisons. Additionally, continuous PM_{2.5} monitoring provides hourly data that assist in understanding how concentrations vary throughout the day. Understanding such daily fluctuations helps determine sources of PM_{2.5} and when health risks from fine particles are greatest. Increased understanding of concentrations and risks aids in prioritizing emission reduction efforts.


PM_{2.5} is a regional pollutant, with some addition from local sources; therefore, concentrations tend to rise and fall in unison across the state. There is considerable variability between sites, however, even as the general trend stays uniform. Such differences in concentrations tend to be driven by local sources, especially those in closer proximity to large urban areas. Disparities between urban and rural areas demonstrate the effect of man-made emission sources on fine particulate concentrations (Figure 4).

Figure 4. Daily average of hourly PM_{2.5} concentrations at several Minnesota sites, weekdays in March 2024



PM_{2.5} emissions in urban areas tend to follow a daily pattern (Figure 5). The mid-morning peak concentration results from traffic. As temperatures rise in the day, the atmospheric mixing height increases. This allows for dilution of fine particle concentrations and lowered concentrations throughout the afternoon. Temperatures fall in the evening, lowering the mixing height and trapping the particles, including those emitted during evening rush hour. This results in elevated concentrations throughout the night.


Figure 5. Average PM_{2.5} concentrations per hour of the day at several Minnesota sites, May 2024



1.3.2. PM₁₀


In 2024, the MPCA currently operated PM₁₀ FRM samplers in St Paul (0866), Minneapolis (0966), and Duluth (0032). The FRM method collects mass samples of PM₁₀ over a 24-hour period once every six days. There are also continuous PM₁₀ FEM monitors that measure hourly PM₁₀ concentrations at sites: St. Paul (0868), Minneapolis (0909), Minneapolis (0910), Minneapolis (1909), Blaine (1002), Marshall (4210), and Virginia (7001) (Figure 6).

Figure 6. 2024 PM₁₀ monitoring sites in Minnesota.



To describe the magnitude of daily PM₁₀ measurements, the MPCA reports the daily PM₁₀ background concentration, which is calculated following the methodology established in EPA’s “PM₁₀ SIP Development Guidance” ([EPA-450/2-86-001, June 1987, Table 6-1](#)). Depending on the total number of samples collected over a three-year period, the daily PM₁₀ background concentration is calculated as the 1st, 2nd, 3rd, or 4th highest daily average PM₁₀ concentration measured over three years. PM₁₀ concentrations ranged from 52 µg/m³ at Minneapolis (0966) to 414 µg/m³ at Virginia (7001) in 2024 (Figure 7).

Figure 7. 2024 PM₁₀ background concentrations compared to the daily NAAQS.



*Site did not meet data completeness criteria

A monitoring site meets the 24-hour PM₁₀ NAAQS when the level of 150 µg/m³ is not exceeded more than once per year, on average, over three years. The standards are attained when the expected number of days per calendar year with a 24-hour average concentration above 150 µg/m³ is equal to or less than one. Minnesota currently meets applicable NAAQS at nine out of ten sites. In 2022-2024, there were no expected exceedances for all sites except Virginia (7001), where the three-year design values was two expected exceedances (Table 2).

Table 2. 2024 PM₁₀ estimated exceedances

Site ID	PM ₁₀ 3-year estimated exceedances	Completeness criteria met
27-003-1002	0	Yes
27-053-0909	0	Yes
27-053-0910	0	Yes
27-053-0966	0	Yes
27-053-1909	0	Yes
27-083-4210	0.3	No
27-123-0866	0	No
27-137-0868	0	Yes
27-137-0032	0	Yes
27-137-7001	2	Yes

1.3.3. Total Suspended Particulate matter (TSP)

TSP was one of the original NAAQS pollutants but was replaced by the PM₁₀ standard in 1987. Generally, smaller particles, such as PM₁₀ and PM_{2.5}, are expected to have greater health impacts than TSP. Today, TSP levels are regulated at the state level by the MAAQS. The MAAQS includes four distinct standards for TSP (Table 3).

Table 3. Minnesota Ambient Air Quality Standards for TSP

Standard type	Time interval	Level of standard	A monitoring site meets the standard if...
Primary ¹	Daily (24-hour)	260 µg/m ³	...the annual 2 nd highest daily TSP concentration is less than or equal to 260 µg/m ³
	Annual	75 µg/m ³	...the annual geometric mean is less than or equal to 75 µg/m ³
Secondary ²	Daily (24-hour)	150 µg/m ³	...the annual 2 nd highest daily TSP concentration is less than or equal to 150 µg/m ³
	Annual	60 µg/m ³	...the annual geometric mean is less than or equal to 60 µg/m ³


¹ primary standard is set to protect against human health effects associated with exposure to an air pollutant.

² secondary standard is set to protect against environmental or public welfare effects associated with exposure to an air pollutant.

In addition to TSP, the filters are analyzed for metals as part of the air toxics program, using the ICP/MS method. Metals are discussed further in the air toxics section of this report.


The MPCA currently operates 21 TSP monitoring sites (Figure 8). Mass samples of TSP are collected over a 24-hour period once every six days.

Figure 8. 2024 TSP monitoring sites in Minnesota




In 2024, Minnesota annual TSP averages ranged from 16 µg/m³ in Blaine (1002) to 44 µg/m³ in St. Paul (0875) (Figure 9). All sites met the annual primary MAAQS in 2024.

Figure 9. 2024 TSP concentrations compared to the annual primary and secondary MAAQS.



In 2024, daily TSP values in Minnesota ranged from 48 $\mu\text{g}/\text{m}^3$ at Bayport (0446) to 192 $\mu\text{g}/\text{m}^3$ at Duluth (7555-1) (Figure 10). St. Paul (0875) and Duluth (7555-1) violated the daily secondary TSP MAAQS; all other sites met the TSP standards.


Figure 10. 2024 TSP concentrations compared to the 24-hour primary and secondary MAAQS.



1.4. Lead (Pb)

The MPCA monitored lead at 21 sites in 2024 (Figure 11), including NCore (1002). Eight lead monitoring sites are considered source-oriented: Gopher Resources Corporation in Eagan (0465), Federal Ammunition in Anoka, MN (6021), Flint Hills (0020 and 0423), North Minneapolis near a former Northern Metals Recycling location (0909 and 0910), Near Road I-94/I-35 (0962), and Bayport (0446), located between two potential sources. The remaining sites are part of the air toxics program.


Figure 11. 2024 lead monitoring sites in Minnesota.



20 out of 21 lead monitoring sites in Minnesota met the 2008 lead NAAQS of $0.15 \mu\text{g}/\text{m}^3$ in 2024 based on the most recent data, which show the three-year maximum 3-month rolling average concentration at monitored sites from 2022-2024. The highest monitored lead design value was $0.46 \mu\text{g}/\text{m}^3$ at Federal Ammunition (6021) followed by $0.14 \mu\text{g}/\text{m}^3$ at Gopher Resources (0465) in Eagan. 18 sites were at levels equal to or less than $0.04 \mu\text{g}/\text{m}^3$ (Figure 12).

The TSP/metals sampler was deployed near Federal Ammunition in Anoka, MN (27-003-6021) in December 2022 to meet the requirement of Appendix D of 40 CFR Part 58, section 4.5. This requirement identifies there must be one source-oriented SLAMS site located to measure the maximum Pb concentration in ambient air resulting from each non-airport Pb source which emits 0.50 or more tons per year. Federal Ammunition identified emissions above the 0.5 tons lead on their annually air emission inventory for 2019 - 2022. The MPCA and EPA are working together on an enforcement investigation to address elevated lead emissions at Federal Ammunition.


Figure 12. 2024 lead design values compared to the 3-month NAAQS.



1.5. Ozone (O₃)


The MPCA monitors ozone at 18 monitoring sites. An additional monitor, located at Voyageurs National Park (0034), is operated by the National Park Service (Figure 13). Since the MPCA does not have any role in the maintenance or use of this monitor, it is not included in our SLAMS or AQI monitoring networks.

Figure 13. 2024 ozone monitoring sites in Minnesota.



A monitoring site meets the primary ozone NAAQS if the three-year average of the fourth highest daily maximum 8-hour concentration is less than or equal to 70 ppb. From 2022-2024, Minnesota eight-hour averages ranged from 53 ppb in Ely (0005) to 66 ppb in St. Michael (3021). All sites were below the eight-hour standard (Figure 14).

Figure 14. 2024 ozone design values compared to the 8-hour average NAAQS.




*Site did not meet data completeness criteria

1.6. Nitrogen Dioxide (NO₂)

Currently, the MPCA monitors NO₂ at six sites in the Minnesota (Figure 15).

Figure 15. 2024 NO_x monitoring sites in Minnesota.



A monitoring site meets the annual NAAQS for NO₂ if the annual average is less than or equal to 53 ppb. Minnesota averages ranged from 4 ppb at Blaine (1002-1), Virginia (7001), and Rosemount (0423) to 9 ppb at the Minneapolis near-road site (0962); therefore, Minnesota met the annual NAAQS for NO₂ (Figure 16).

In addition to the annual standard, there is also a one-hour standard for NO₂. The one-hour NAAQS is intended to protect against adverse health effects associated with short-term exposures to elevated NO₂. To meet this standard, the three-year average of the annual 98th percentile daily maximum one-hour NO₂ concentration must not exceed 100 ppb. Minnesota averages ranged from 24 ppb at Rosemount (0423) to 43 ppb at Minneapolis (0962); therefore, all sites met the one-hour NAAQS for NO₂ (Figure 17).

Figure 16. 2024 NO₂ design values compared to the annual NAAQS.






Figure 17. 2024 NO₂ design values compared to the 1-hour NAAQS.



1.7. Carbon monoxide (CO)

The MPCA monitors CO at six sites in Minnesota, all in the Twin Cities metropolitan area (Figure 18).

Figure 18. 2024 CO monitoring sites in Minnesota.



Minnesota currently meets applicable eight-hour NAAQS for CO. A monitoring site meets the 8-hour CO NAAQS when the level of 9 ppm is not exceeded more than once per year. A design value is a statistic that describes the air quality status of a given location relative to the level of the National Ambient Air Quality Standards (NAAQS). The design value for CO is evaluated over a two-year period. Specifically, the design value is the higher of each year's annual second maximum, non-overlapping 8-hour average. Minnesota CO design values for 2024 for ranged from 0.8 ppm at Blaine (1002) to 3.8 ppm at Minneapolis (0954) (Figure 19).

The one-hour CO NAAQS is met when the level of 35 ppm is not exceeded more than once per year. A design value is a statistic that describes the air quality status of a given location relative to the level of the National Ambient Air Quality Standards (NAAQS). The design value for CO is evaluated over a two-year period. Specifically, the design value is the higher of each year's annual second maximum, non-overlapping 1-hour average. Minnesota values for 2024 ranged from 1.1 ppm at Rosemount (0020) to 11.7 ppm at Minneapolis (0954) (Figure 20).

Figure 19. 2024 design values for CO compared to the 8-hour NAAQS.






Figure 20. 2024 CO design values compared to the 1-hour average NAAQS.



1.8. Sulfur dioxide (SO₂)


The MPCA monitors SO₂ at six sites in the Twin Cities metropolitan area (Figure 21) and one site in Virginia (7001). Trace level SO₂ at the NCore site in Blaine (1002) will help us understand the role of SO₂ at levels far below the NAAQS.

Figure 21. 2024 SO₂ monitoring sites in Minnesota.



The primary SO₂ NAAQS is a one-hour standard; it is met if the three-year average of the annual 99th percentile daily maximum one-hour SO₂ concentration is less than 75 ppb. Minnesota averages from 2022-2024 ranged from 2 ppb at Blaine (1002) to 14 ppb at Rosemount (0020); therefore, all Minnesota sites met the one-hour NAAQS for SO₂ (Figure 22).

Figure 22. 2024 SO₂ design values compared to the 1-hour NAAQS.



*Site did not meet data completeness criteria

2.AQI

The AQI categories developed by the EPA are green (good), yellow (moderate), orange (unhealthy for sensitive groups or USG), red (unhealthy), purple (very unhealthy), and maroon (hazardous). Each category is assigned a color and corresponds to a different level of health concern (Figure 23). In the past, MPCA offered AQI forecasts for only the Twin Cities and Rochester. The MPCA issues daily forecasts for the majority of its AQI monitor locations through MPCA’s AQI website, the Minnesota Air mobile app, Twitter, and to individuals who have signed up to receive e-mailed forecasts. If it is suspected through forecasting or monitoring that the daily AQI will be over 100, the MPCA will issue an Air Quality Alert to be disseminated by the National Weather Service, GovDelivery, the AirNow mobile app, Twitter, the media, and to individuals who have signed up to receive email alerts. Forecasts and alerts allow the public to be proactive about protecting their health and reducing their own contributions to emissions and exposure to air pollution. At no cost, the public can download the Minnesota Air mobile app and sign up for emailed forecasts and alerts from [MPCA’s AQI website \(https://www.pca.state.mn.us/air/current-air-quality\)](https://www.pca.state.mn.us/air/current-air-quality).

Figure 23. The Air Quality Index categories and respective levels of health concern.

Air Quality Index Levels of Health Concern	Numerical Value	Meaning
Good	0 to 50	Air quality is considered satisfactory and air pollution poses little or no risk.
Moderate	51 to 100	Air quality is acceptable; however, for some pollutants there may be a moderate health concern for a very small number of people who are unusually sensitive to air pollution.
Unhealthy for Sensitive Groups	101 to 150	Members of sensitive groups may experience health effects. The general public is not likely to be affected.
Unhealthy	151 to 200	Everyone may begin to experience health effects; members of sensitive groups may experience more serious health effects.
Very Unhealthy	201 to 300	Health alert; everyone may experience more serious health effects.
Hazardous	301 to 500	Health warnings of emergency conditions. The entire population is more likely to be affected.

Compared to 2023, Minnesota experienced a quieter year with respect to Orange or worse AQI days for PM_{2.5} during 2024 (Figure 24). Above normal precipitation during the spring helped suppress local fire activity, and relatively wetter conditions north of the border led to milder fire season in western and eastern Canada. There were 7 days with an AQI of orange or worse in 2024, which was near average. Northern portions of the state experienced more periods of smoke (Figure 25). The vast majority of days in 2024 were Green at any single monitor.

Figure 26 shows the maximum daily PM_{2.5} AQI throughout Minnesota in 2024. Yellow AQI winter stagnation episodes occurred in January and February, and again in November and December. Prescribed fire led to a few marginal yellow days during the spring. The worst air quality of the year was associated with several intrusions of wildfire smoke from Canada. AQIs reached Red during smoke intrusions in mid-May and mid-September and reached Orange on a few days between mid-August and early September.

Figure 24. Annual statewide PM_{2.5} AQI category counts, 2001-2024.






Figure 25. 2024 daily observed AQI category counts for PM_{2.5} in Minnesota.



*Voyageurs N.P. a non-forecast location

Figure 26. Peak daily PM_{2.5} AQI for Minnesota in 2024.



AQIs with respect to ozone were fairly benign in 2024 (Figure 10). The spatial extent of ozone followed the typical pattern, with higher ozone tending to be found near and downwind of the Twin Cities (Figure 26). The weather in Minnesota over the spring and early summer of 2024 was cloudy and wet, conditions which typically suppress ozone. The only Orange AQI occurred in late July due to pollution pooling along a weak frontal boundary (Figure 27). The majority of Yellow-Orange days occurred from mid-April to the end of July, which is typical for Minnesota (Figure 28).

Figure 27. Annual statewide ozone AQI category counts, 2001-2024.






Figure 28. 2024 daily observed AQI category counts for ozone in Minnesota (during the official ozone season from March 1 to October 31).



*Voyageurs N.P. a non-forecast location

Figure 29. Peak daily ozone AQI for Minnesota in 2024.



The MPCA issued a six air quality alerts covering 14 days during 2024 (Figures 30 and 31). All of the alerts in 2024 were due to wildfire smoke.

Figure 30. Total count and causes of air quality alert forecasts issued in Minnesota counties, 2010-2024.





Figure 31. Total count and causes of air quality forecast alert days issued in Minnesota counties, 2010-2024.



3. Air Toxics

Air toxics are routinely monitored in Minnesota as part of the MPCA's ambient air toxics monitoring program. The MPCA monitors three types of air toxics: VOCs, carbonyls, and metals. To determine the health risk of the measured air toxics concentrations, the MPCA uses guidelines, called inhalation health benchmarks.

A compiled list of the health benchmarks can be found in the [Ambient air toxicity values workbook \(https://data.pca.state.mn.us/views/Airtoxicityvalues/Airtoxicityvalues?%3Aembed=y&%3AisGuestRedirectFromVizportal=y\)](https://data.pca.state.mn.us/views/Airtoxicityvalues/Airtoxicityvalues?%3Aembed=y&%3AisGuestRedirectFromVizportal=y).

These health benchmarks come from a variety of sources, including the following:

- [Minnesota Department of Health air risk guidance \(https://www.health.state.mn.us/communities/environment/risk/guidance/air/table.html\)](https://www.health.state.mn.us/communities/environment/risk/guidance/air/table.html)
- [EPA's Integrated Risk Information System \(IRIS\) \(https://www.epa.gov/iris\)](https://www.epa.gov/iris)
- [California's Office of Health Hazard Assessment \(OEHHA\) \(https://www.oehha.ca.gov/air.html\)](https://www.oehha.ca.gov/air.html)
- [EPA's Superfund Program \(https://www.epa.gov/superfund\)](https://www.epa.gov/superfund)

MPCA is currently in the process of updating the state-wide air toxics monitoring results.

There are select monitoring sites where measured concentrations are shared publicly. Data explorers for the following sites are available on [MPCA's Data visualizations webpage \(https://www.pca.state.mn.us/about-mPCA/mpca-data-visualizations\)](https://www.pca.state.mn.us/about-mPCA/mpca-data-visualizations).

- Metals at Saint Paul west side community
- Lead at Federal Ammunition
- Air toxics at Otter Lake Technologies

4. Visibility


Air pollution can reduce visibility over wide areas, a phenomenon called regional haze. Haze occurs when sunlight encounters fine particles in the air, which absorb and scatter light. Haze-causing pollutants come from a variety of sources, both natural and human-made, including motor vehicles, electric utilities, taconite processing facilities, agricultural activities, and wildfires.

In 1999, the EPA established a regulatory program to reduce haze caused by human-made air pollution at national parks and wilderness (Class I) areas. The goal of the regional haze rule is to achieve natural visibility conditions in Class I areas by 2064, with interim progress goals set every 10 years. The first interim progress goal was set for 2018.

Minnesota has two Class I areas, both located in Northern Minnesota—the Boundary Waters Canoe Area Wilderness and Voyageurs National Park (Figure 32).

Visibility is calculated from PM_{2.5} species measurements through the [IMPROVE Aerosol Network website \(http://vista.cira.colostate.edu/IMPROVE/\)](http://vista.cira.colostate.edu/IMPROVE/). Figure 14 shows the location of Minnesota sites in the network. Minnesota has an IMPROVE site in each of the two Class I areas. An additional site was installed at the Great River Bluffs State Park in southeastern Minnesota to help better understand the regional transport of pollutants that impair visibility into Minnesota from the Southeast.

Figure 32. Class I areas in Minnesota impacted by regional haze.



The IMPROVE network PM_{2.5} speciation measurements are mathematically processed to express visibility as a five-year rolling average deciview (dv) value. A human observer is thought to be able to visually perceive a one to two deciview difference in scene appearance. The MPCA aims to see calculated deciview values on the most impaired visibility days reach natural conditions by 2064. Interim goals are set for every ten years. Both the Boundary Waters and Voyageurs sites achieved the 2018 interim goal by 2012 and reached the 2028 interim goal in 2023. Visibility on the clearest days at both sites has not degraded over time and have actually improved (Figure 33).

Figure 33. Visibility progress measured at Boundary Waters and Voyageurs (2004-2023) and the estimated interim progress goals for 2018 and 2028.

