MINNESOTA POLLUTION CONTROL AGENCY AQUATIC LIFE CRITERIA AND WATER QUALITY STANDARDS

Page 1 SUMMARY

•	A. Chemical/Element:	CAS# 1763231	Dates Proposed/Promulgated	Aquatic Tox. &
_	Perfluorooctanesulfonic acid (PFOS)		/	Human Health
Arc.	Site-specific Pool 2, Mississippi R.		Developed 3Aug2007	AL and HH
1	i _e		Revised 30 Oct 2009	НН
	(V)		Revised 23 January 2013	HH
•	CU			

	sota Water Quality Standards:	μg/l (unless noted	d otherwise)		
Water Class 1, 1,2A	Use	CS	MS	FAV	CS Basis ¹
1,2A	DW, Salmonid	14 ng/L	85 μg/L	170 μg/L	PCA Hs
1,2Bd	W W, NonSalmonid	14 ng/L	85 μg/L	170 μg/L	PCA Hs
2B, 2C, 2D	Non Salmonid	14 ng/L	85 μg/L	170 μg/L	PCA Hs
	Other 1				
CS: Chronic Sta	andard, DW: G rinking Water	, FAV: Final Acute	Value, MS: Ma	ximum Standard	
Toxicity related	andard, DW: Spinking Water				
If yes, above cri	iteria values determine tor:				
Slope: Acute:	: Q _{UQ}), :: MPCA				
Chronic	e: ~//	Ö.			
Formulas:	MPCA	, C.		EPA	

Formulas:	MPCA S _x	EPA
CS:	And.	
MS:	Arde	
FAV:	°Uni	

Notes:

		·		
C.	EPA Criterion: μg/l	CCC: none	Basis:	
	Date:	MC: none	Basis:	
		FAV: none	B ć is:	

C. EPA Criterion: µg/1	ccc: none	Basis:	
Date:	MC: none	Rasis:	
	FAV: none	Brois:	
		784,	
D. Other Criteria μg/l		Ny Source	
0.2 (based on 10 kg child an	d 1 L drinking water	EPA Office of Water Provisional Health	
intake)	_	Advisory under the safe Drinking Water	
		Program at	
		http://www.epa.gov/waterscience/criteria/drin	
		king/	
E. Notes: Mississippi River site-s	pecific criterion for PFOS	is based on BAF information collected from Pool 2.	
1		EPA Office of Water Provisional Health Advisory under the Safe Drinking Water Program at http://www.epa.gov/waterscience/criteria/drin king/ is based on BAF information collected from Pool 2, iterion gency staff res used	
Criteria basis codes for part B:	(55.)		
EPA = From U. S. Environmental	Protection Agency (EPA) cr	iterion	
PCA = Criterion developed by M	innesota Pollution Control Ag	gency staff	
T1 = Direct aquatic life toxicity, I	EPA national criteria procedu	res used	٥.
T2 = Direct aquatic life toxicity, I	EPA advisory procedures used	d	1
Hs = Human health systemic effe	cts		1/0/2
Hc = Human health carcinogenic	effects		100
R = Tissue residue (bioaccumulat	ion)		Ÿ,
W = Wildlife effects			

O = Organoleptic (taste and odor)

Other = Criterion based on other end point

MINNESOTA POLLUTION CONTROL AGENCY AQUATIC LIFE CRITERIA AND WATER QUALITY STANDARDS

Page 2 DIRECT AQUATIC LIFE TOXICITY - EPA Criterion Available

A. Ch	emical/Element	: PFOS		CAS# 1763231			
						_	
В.	EPA Criterion	n: µg/l	CCC:	none		Basis:	
120	Date:		MC:	none		Basis:	
90			FAV:	none		Basis:	
~~~	42						
1.	Related to wa	ter quality?: no					
		1=			1		1
2.	Toxicio	FAV:			N:		ACR:
	~ (/	ter quality?: no  FAV: Chronic value:			N:		
3.	Residue	Dru Administratio					
	Food and I	Drug Administration	on (FDA	) action level:			
	BCF Final	l: Contact 1% lipid	N total:	N used:			
	geo mean a	at 1% lipid					
	% lipid:	'h					
	geo man u	nadjusted for lip	<u> </u>				
ACR:	Acute to Chron	ic Ratio, BCF: Bio	ocenti	ration Factor, CCC	: Chroni	c Criterion, l	MC: Maximum Criterion
_			1/2	•			
C.	MPCA Evalua	ation of EPA Crite	erion of				
	1. Four lowes	st GMAVs (Genus	Mean A	coge values):			
	2. Commerci	ally or recreationa	lly impo	N used:  ration Factor, CCC  Report Values):  rtant species:			
	0 DI . 1 .			"For			
	3. Plant data:			no			
	4 Extrapolat	ion of water qualit	y offoots	. ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	in.		
	4. Extrapolat	ion of water qualit	y effects	•	Torn		
	5. Chronic da	ata No. of	values:		Pati		
	5. 224341 <b>2 0</b> 0	No. bel	ow criter	rion:	-0	<b>7</b>	
						The	
	Notes:					. 100°.	
						"hy	,
	6. ACRS	ACR used by EI	PA:		N:	-4	<i>y</i>
	1	10 11 4	CD.				· -/ 1.

6. ACRS	ACR used by EPA:	N:	
	Geo. mean, all ACRs:	N:	
	ACR used by MPCA: 9.12	N: 3	
Separate Coo No. of Sa N(nonsal) Adjustme	has no surface water criteria for PFOS  I/Warm Water Criterion, ug/l Imonids deleted from lowest 4 GMAVs: D: FAV: MC: Ents to FAV:	CC:	Nater quality standards.
Notes:	1 DDA 1		(any
Summary of o	changes made to EPA criterion		Sards,

D. Separate Cool/Warm Water Criterion, ug/l

E. Summary of changes made to EPA criterion

## Page 3 DIRECT AQUATIC LIFE TOXICITY when no EPA criterion is available

A	. Chemical/I	Element:	PFOS	CAS	# 17632	31	
В.	EPA N	Jational M	Iethod Salmonid (2A water only) Osteichthyes (fish): Chordata (fish, amphibian Planktonic crustacean: Benthic crustacean: Aquatic insect: Phylum other than Arthropy Second insect or phylum (2A): See Tier II method (2A): See Tier II method (3A):  Table 2a  No.  Acute value 9100 67,200 Generic				
VIL.	Data require	ements:	Salmonid (2A water only	):		Onchoryn	chus mykiss
<del>α</del> ζ	· ·		Osteichthyes (fish):	<i>)</i> -		Pimephale	es promelas
	*OC//.		Chordata (fish, amphibiar	n):		Pseudacri	1
	TO TO THE PARTY OF		Planktonic crustacean:			Daphnia n	v
	To.		Benthic crustacean:			Hyalella a	
	7/	De	Aquatic insect:				-
		Orn	Phylum other than Arthro	poda or C	hordata:	Unio comp	plamatus
		Price	Second insect or phylum	not already	v rep.:	Lumbricul	lus variegatus
2.	GMAVs	Lowest	(2A): See Tier II method	1	Lowes	t 4(2B,2C, 2I	D): See Tier II method
	μg/l		7/2				,
		N:	Cox		N:		
			Tac.				
3.	FAV:	2A:	W.		2B, 2C	, 2D:	
4.	Adjustment	s to FAVs	S: Yer		•		
5.	Chronic c	lata: Se	e Table 2a No.		S	pecies:	
	mean valı	ues	7/3				
	μg/l		U.S.				
	<u> </u>		70	γ _ο ,			
6.	ACR Meas	ured:	Acute value	%, Ch	ronic val	ue	ACR
			9100	741	0		22.19
G	eneric: 18		67,200	36	350		1.9
			Generic	Ge	n <b>es</b> ic		18
Fi	nal: 9.12				'eip		
					70,	<b>%</b>	
	Final Plant		NOEC = $300 \mu g/L$ , Northe	ern milfoil		Day.	
8.	Chronic Cri	iterion (FA	AV/ACR) see Tier II	method		0/2	-
						41 hr	1.9 18
C.	EPA A	Advisory N	Method (Tier II method)			<i>'D</i> ₃	·
1	Doto roquir	amanta	Eig	h.		Dimanhal	Manamalas

C. EPA Advisory Method (Tier II method	d)	<i>'D</i> y.
1. Data requirements:	Fish:	Pimephales <b>'s</b> romelas
N = 7	Crustacean:	Daphnia magi <b>li</b>
No. SMAVs: 8	Third animal:	Lumbriculus varies atus
No. GMAVs: 7	Plant for herbicide:	`\$\f\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
Adjustment Factor: 4.3	Insect for pesticide:	```.'/Ŋ _k

Adjustment Factor: 4.3 Insect for pesticide:

2. Lowest GMAV: 5600 μg/L Species: Lumbriculus variegatus

3. FAV: 1302 μg/L (reduced to 170 μg/L) MC: 85 μg/L

4. Chronic data: See B.5.

5. ACR: 9.12 (See B.6)

6. CC: 18.6 μg/L (rounded up to 19 μg/L)

7. Citation for lowest GMAV: STS-334

D. Notes: FAV = GMAV ÷ Adjustment Factor; EAO staff lowered the Tier II calculated FAV of 1302 μg/L to the project EC50 of 170 μg/L to protect Chironomus tentans. The original Chironomus tentans study used the highest exposure concentration of 150 μg/L and reported the EC50 as greater than 150 μg/L. Therefore, EAO staff determined an estimated EC50 for the test at 170 μg/L.

## MINNESOTA POLLUTION CONTROL AGENCY AQUATIC LIFE CRITERIA

#### Page 4 HUMAN HEALTH

Source: MDH   Source: MDH	A. Chemical/Element: PFOS	CA	<b>AS#</b> 1	1763231			
Reference Dose:   mg/kg/day   Cancer Potency Slope:   (mg/kg-d)¹		DW 16	1	C' 1 1		DW 1	D. Cl
Source: MDH   Source: MDH	B. EPA Human Health Criterion (μg/l):		sn:	fish only	none	-	
Source: MDH   Source: MDH	Le _O						
Source: MDH   Source: MDH	" 40 _{C1.}						e 0.2 μg/L
Relative Source Contribution (RSC):    C.   Minnesota man Health Criterion	Reference Dose.		tency	Slope:			$(mg/kg-d)^{-1}$
C. Minnesta imman Health Criterion  1. Ref.dose: 0.0008	Final BCF.	%lipid:					
Neef.dose: 0.00008	Relative Sources Contribution (RSC):						
1. Ref.dose: 0.00008	C. Minnesota Wanner Health Critarian						
Additivity endpoint(s): Evelopment (body weight/weight gain), Hepath (liver) system, Thyroid  RSC: 0.2  2. Cancer Potency Slope: n/a  3. Measured BAFs: Species/Tissue  4. Bluegill/ Fillet  5. Carp/ Fillet  6. Smallmouth Bass/ Fillet  7. Smallmouth Bass/ Fillet  7. Edible portion BAF or BCF  Cold water: 6.0 % lipid  7. log Kow:  adjust. for % lipid:  8. Parachor: n/a  9. BCF to BAF conversion factor: n/a  10. Final BAF: 2A (6% lipid): 2667  11. Criteria:  and conversion factor: n/a  Source: MDH	1 Paf dosa: 0.000082	ma/ka/day	So	urca: MD	Ц		
RSC: 0.2   Source: MDH	Additivity and point(s): Rayalanment (body	mg/kg/uay	_				
RSC: 0.2   Source: MDH		Thyroid	30	uice. Mid	11		
1.	RSC: 0.2	111/1010	So	urce: MD	H		
2. Carp/ Fillet 1237 n/a 1237 3. Freshwater Drum / Fillet 3077 n/a 3077 4. Smallmouth Bass/ Fillet 2845 n/a 2845 5. White Bass/ Fillet 3078 d618 n/a 4618  Geo mean: 2667 4. Measured BCFs: Species/Tissue 2667 4. Measured BCFs: Species/Tissue 1078 none 1078  Geo mean: 1078  Seo mean: 1078  Geo mean: 1078  Seo mean unadjusted for lipid: 1078  Seo mean: 1078  Seo m	2. Cancer Potency Slope: n/a (m	ng/kg-d) ⁻¹	_		-		
2. Carp/ Fillet 1237 n/a 1237 3. Freshwater Drum / Fillet 3077 n/a 3077 4. Smallmouth Bass/ Fillet 2845 n/a 2845 5. White Bass/ Fillet 3078 d618 n/a 4618  Geo mean: 2667 4. Measured BCFs: Species/Tissue 2667 4. Measured BCFs: Species/Tissue 1078 none 1078  Geo mean: 1078  Seo mean: 1078  Geo mean: 1078  Seo mean unadjusted for lipid: 1078  Seo mean: 1078  Seo m	3. Measured BAFs: Species/Tissue				9	6 lipid	Norm BAF
1.	1. Bluegill/ Fillet <b>Q</b>		27	00			
1.	2. Carp/Fillet	,	12:	37	n/a		1237
1.	3. Freshwater Drum / Fillet	S _E	30	77	n/a		3077
1.	4. Smallmouth Bass/ Fillet	<i>nd</i> 3.	28	45	n/a		
1.	5. White Bass/ Fillet	70%	46	18	n/a		4618
The second color of the		Un,	<b>je</b>				
1.	Geo mean:		<b>6</b> 26	67			
The second color of the	4. Measured BCFs: Species/Tissue		7	BCF	9	6lipid	Norm. BCF
8. Parachor: n/a  9. BCF to BAF conversion factor: n/a  10. Final BAF: 2A (6% lipid): 2667  11. Criteria: 2A: 14 ng/L * 2Bd: 14 ng/L * 2B/2C, 2D: 14 ng/L # HRL/HBV: 0.3 μg/L MDH Health Risk Limit/Health Risk L	1.		no	ń <b>€</b>			
8. Parachor: n/a  9. BCF to BAF conversion factor: n/a  10. Final BAF: 2A (6% lipid): 2667  11. Criteria: 2A: 14 ng/L * 2Bd: 14 ng/L * 2B/2C, 2D: 14 ng/L # HRL/HBV: 0.3 μg/L MDH Health Risk Limit/Health Risk L			-	DATE			D.CE
8. Parachor: n/a  9. BCF to BAF conversion factor: n/a  10. Final BAF: 2A (6% lipid): 2667  11. Criteria: 2A: 14 ng/L * 2Bd: 14 ng/L * 2B/2C, 2D: 14 ng/L # HRL/HBV: 0.3 μg/L MDH Health Risk Limit/Health Risk L			/-	BAF 'A	<u>.</u> 2.		BCF
8. Parachor: n/a  9. BCF to BAF conversion factor: n/a  10. Final BAF: 2A (6% lipid): 2667  11. Criteria: 2A: 14 ng/L * 2Bd: 14 ng/L * 2B/2C, 2D: 14 ng/L # HRL/HBV: 0.3 μg/L MDH Health Risk Limit/Health Risk L			n/a	<u>'</u>	9× .		
8. Parachor: n/a  9. BCF to BAF conversion factor: n/a  10. Final BAF: 2A (6% lipid): 2667  11. Criteria: 2A: 14 ng/L * 2Bd: 14 ng/L * 2B/2C, 2D: 14 ng/L # HRL/HBV: 0.3 μg/L MDH Health Risk Limit/Health Risk L	*		n/2	<u> </u>	· http		
8. Parachor: n/a  9. BCF to BAF conversion factor: n/a  10. Final BAF: 2A (6% lipid): 2667  11. Criteria: 2A: 14 ng/L * 2Bd: 14 ng/L * 2B/2C, 2D: 14 ng/L # HRL/HBV: 0.3 μg/L MDH Health Risk Limit/Health Risk L	, <u> </u>		me	i vac	OSAR		Fet BCF:
8. Parachor: n/a  9. BCF to BAF conversion factor: n/a  10. Final BAF: 2A (6% lipid): 2667  11. Criteria: 2A: 14 ng/L * 2Bd: 14 ng/L * 2B/2C, 2D: 14 ng/L # HRL/HBV: 0.3 μg/L MDH Health Risk Limit/Health Risk L			1110	as.	lipid):	Why.	Lst. BCI .
10. Final BAF: 2A (6% lipid): 2667  11. Criteria: 2A: 14 ng/L * 2Bd: 14 ng/L * 2B/2C, 2D: 14 ng/L # HRL/HBV: 0.3 μg/L MDH Health Risk Limit/Health Risk Lim	·				<b>F</b> /-	<i>'</i> O ₂	
11. Criteria: 2A: 14 ng/L * 2Bd: 14 ng/L * 2B/2C, 2D: 14 ng/L # HRL/HBV: 0.3 μg/λ MDH Health Risk Limit Heal						ک ^ب ن	ريار الاي
11. Criteria: 2A: 14 ng/L * 2Bd: 14 ng/L * 2B/2C, 2D: 14 ng/L # HRL/HBV: 0.3 μg/λ MDH Health Risk Limit Heal	10. Final BAF: 2A (6% lipid): 2667	2B &	2Bd,	2C, 2D (1.	5% lipio	1): 2667	16.12
D 137.1 C C 1 <b>9%</b>	11. Criteria: 2A: 14 ng/L * 2Bd: 14 ng/L	* 2B/20	C, 2D	: 14 ng/L	# HR	L/HBV: 0	.3 μg <b>λ</b>
Based Value for Ground Water	ug/l				MD	H Health I	Risk Link Health
					Bas	ed Value for	or Groundwater
	D 0 1 .: /			<u> </u>			ed in Minn R. ch 705
D. Organoleptic: n/a Source:	D. Organoleptic: n/a		2	source:			
CII. I VJV.	Ch. 7050.  Data used for calculation of fish BAF value	es for PFOS	S Wer	e renorte	l fish ti	ssue and s	urface water sami
Data used for calculation of fish BAF values for PFOS were reported fish tissue and surface water s	pata used for calculation of fish DAT value				4 11811 (1) 11	souc and s	arrate water sample

collected from the Mississippi R, Pool 2 in 2012. Methods used for calculating water quality criteria can be found in the MPCA water quality guidance manual (Maschwitz, 2000).

#### Water Quality Criteria development for 2012 Mississippi River PFOS sample analysis

Individual fish tissue values were measured from fillets of fish captured from pool 2 in 2012 and from surface water samples collected from the same area (Table 1). Procedures and results of those measurements are reported elsewhere. Briefly, fish and water were collected from the sites throughout pool 2 segregated by four discrete sampling sections of the pool. An arithmetic average of the measured water concentrations was calculated for case. The pool of the pool of the measured tissue value divided by the average measured tissue value divided by the average measured to the fish collection site. These individual BAFs were used to calculate a pool-wide geometric mean BAF. Values analyzed and reported as non-detect were given the value of the reporting final for that analysis. Presently, guidance is being developed to better serve MPCA in its use of censored data.

1/2				Species BAF (L		ı	
ion on		Blue Gill Sunfish	Carp	Freshwater	Smallmouth	White	Section
100	<b>/</b> _1		1210.32	Drum 6106.25	Bass 5092.91	Bass 8726.25	GeoMean 4138.21
u	£0.	4064.24	052.26	2526.50	1922.07	0166.20	27/0 21
<u>i</u>	3 804	4304.34	1605.13	3060.38	3608.00	5627 30	4405.58
×	1	4383.94 598.29	/132.09	10/15 7	738.82	882 12	702.35
		<b>8</b>	+32.03	1043.7	Grand Geom	etric mean	2632.09
measured easured fr	from fish om the co	orresponding	sach section.	n divided by th	e arithmetic ave	rage water c	oncentration
fCC =	nd water	RfD x BW x I	RSC	Thoreing			
Where:	fCC = f RfD = r BW = s RSC = exposu consun IW = in CR = fi BAF =	w + (BAF x) ish consumption reference dose standard body Relative Source re fraction attrination (0.2) cidental ingesish consumption Biological Acc	on criterion $e = 0.00008$ weight (70 less Contributed to watten of water on rate (0.03) $control = 0.000$	(mg/L) (mg/kg-d) kg) tion factor: ater and fish r (0.01 L/d) 80 kg/d) Factor (L/kg)	3608.00 738.82  Grand Geometric average arithmetic	*de.nn.lishar	
0.01	2632.09 <i>L,</i> sh tissue 2.09	BAFs for all	of Pool 2:	o,		<b>.</b> 67	Mater quality st

$$\frac{RfD \times BW \times RSC}{[IW + (BAF \times CR)]}$$

$$fCC = \frac{0.00008mg/kg - d \times 70 \ kg \times 0.2}{[0.01 \ L/d + (2632.09 \ L/kg \times 0.030 \ kg/d)]} = 0.0000142 \ mg/kg$$

$$BAF (L/Kq) = 2632.09$$

$$fCC (ng/L) = 14$$